

HumaCount 5D

Outstanding 5-part Diff Hematology System

- > Direct Capillary Process by OptimalCount Technology
- > Distinct 5-part diff
- > Definite Immature Cell Count (LIC, ALY)

CoreLab DX

HEMATOLOGY



Human

Diagnostics Worldwide

5-part Diff

Importance of white blood cell differential count


Benefits of a 5-part diff

- > Better, targeted assessment of the immune response
- > Reduced number of manual blood smears
- > Faster reporting time
- > Saves costs

*«A 5-part diff is essential for
Eosinophils (EOS)/
Neutrophils (NEU) determination»*

5-part diff provides a clear picture of the immune status

The WBC differential divides the white blood cells into the 5 major sub-populations. Each cell type provides information about an immune response or a disease type



Importance of EOS and NEU separation for a targeted diagnosis

- > A high number of eosinophils EOS indicates a parasitic infection
- > A high number of neutrophils indicates a bacterial infection

A 3-part system groups cell types, hence providing only limited information on the disease status

- > **MID** = MON + EOS
- > **GRA** = NEU + EOS + BAS

For the full picture manual blood smears are required for a high performance or an outstanding 5-part diff system like HumaCount 5D.

HumaCount 5D

Innovations you can count on

5-part diff hematology analyzer

- > Small footprint stand-alone system with integrated PC
- > 29 parameters with ATL#% & LIC#%
- > Sample volume: 20 μ l
- > Up to 60 samples / hour
- > 2D barcode target value transfer



Direct capillary blood process using OptimalCount Technology

Same accuracy compared to venous blood. Total capillary blood volume of 20 μ l, 0 μ l dead volume.

Distinct 5-part diff

Providing excellent differentiation of NEU, EOS, MON, LYM and BAS, based on 3D scatter technology.

Definite immature cell count

Quantitative count and percentage value of Large Immature Cells (LIC), Atypical Lymphocytes (ALY).

5-part diff or CBC-count mode switch

One click option to switch between full 5-part diff or CBC-count for each sample.

Direct Capillary Blood Process


Easy, less painful 5-part diff results from one drop of blood

OptimalCount Technology for capillary samples


- > Accuracy as exact as for venous samples
- > Blood volume defined by capillary tube
- > Total sample volume 20 μ l, 0 μ l dead volume
- > Dilution defined by auto-diluent dispensing
- > No manual steps needed



Direct capillary blood process with OptimalCount Technology



Conventional capillary mode – error prone manual method




«OptimalCount Technology guarantees, accuracy as exact as with venous samples, 20 μ l sample volume, 0 μ l dead volume, and precise dilution – thanks to auto-dispensing.»



Benefit of Capillary Blood Sample


- > No physician needed to collect capillary blood
- > Quick, simple and less painful blood collection
- > Especially required for infants and small children, elderly with fragile veins and severely burned patients
- > Equally suitable for use with children and adults

Defined high cell count rate



Correct dilution ratios due to auto-diluent suspension, blood volume defined by capillary tube, plus a count of ~3000 cells result in high accuracy normally only possible with venous samples

Defined dilution ratio (process) auto-aspiration of partially diluted sample



Many error-prone manual steps result in incorrect dilution ratios. A very low number of cells counted in a pre-diluted sample leads to very low accuracy with conventional analyzers.

Dilution inside analyzer

Distinct 5-part Diff

Improved clinical utility

Target diagnosis and treatment with 5-part diff


- > Absolute count and percentage of each parameter, NEU, EOS, MON, BAS, LYM, with immediate clinical relevance
- > Ability to detect abnormal cells, LIC, ALY
- > Overcomes restrictions of 3-part systems such as grouping of cell types such as MON/EOS and NEU/MON/BAS

Better differentiation with 3D scatter technology

3-channel laser detection channel (3D) for:


- > Eosinophils (EOS)
- > Neutrophils (NEU)
- > Monocytes (MON)
- > Lymphocytes (LYM)
- > Basophils (BAS)

Scatter diagram with all parameters




5-part diff and LIC, ALY parameters


3-channel laser detection for EOS, NEU, MON, LYM



Scatter angle 1




Scatter angle 2




Scatter angle 3

BAS detection channel



Dedicated BAS detection channel



NORMAL CONDITION

Definite Immature Cell Count

Reliable immune cell analysis (LIC, ALY)

LIC and ALY without blood smears

3D Laser Scatter enables quantitative count and percentage value of Large Immature Cells (LIC) and of Atypical Lymphocytes (ALY).


LIC (blasts)

are an excellent routine parameter indicating the balance between leucocyte production, circulation in the body and consumption due to immune defense. A high number of LICs is often described as a 'left-shifted' leukogram, with more banded cells and metamyelocytes present while segmented neutrophils are already depleted in the blood.

ALY (lymphoblasts)

are larger than naive lymphocytes.

ALY are lymphocytes that have increased in size due to activation by an antigen, which in turn triggers increased synthesis of mRNA and protein. ALY in blood are always an alarm signal. ALY are seen in the blood of patients suffering from acute lymphoblastic leukemia (ALL); viral disease such as cytomegalovirus, Epstein Barr Virus, hepatitis C; bacterial infections such as toxoplasmosis; exposure to radiation; drug and immunization reactions; and other immune responses.




5-part Diff or CBC-count Mode Switch

Flexible and efficient with one click

One-click reagent optimization

Not every patient sample requires 5-part diff

- > Increase your flexibility with our one click option
- > Switch between full 5-part diff and CBC-count for each sample
- > Optimize your costs by reducing your reagent consumption
5-part diff = 3 reagents / CBC-count = 2 reagents



One-click sample recording

STAT samples require fast action

A new sample is recorded with a one-hand operation. When the sample is positioned under the needle for aspiration, the recording of parameters is started with the same hand by depressing the large red switch. Automated print-out and data transfer via LIS are supported.

Intuitive, ICON based HUMAN software

- > One click to result
- > One screen provides an overview of all 29 parameters, scatter plots and flags

Different tube types supported

- > Small and large EDTA primary tubes
- > Bullet tubes / capillary tubes

HumaCount 5D System Reagents

Reagents* REF

HC5D-Diluent _____ 16450/10

- > Contains 20 l

HC5D CBC-Lyse _____ 16450/20

- > Contains 200 ml

HC5D-Diff-Lyse _____ 16450/30

- > Contains 500 ml

HC5D-Clean _____ 16450/60

- > Contains 50 ml

* require RF card

Control REF

HC5D Control _____ 16450/40

- > Target value upload via 2D barcode
- > 3 Levels, multi-parameter
- > 2 x 3 x 3 ml

Calibrator

HC-Calibrator _____ 17400/50

- > For use on all HUMAN hematology systems
- > 1 x 2 ml



981646/2018-03 © 2018 Human

